

COOL AND HOT COMPONENTS OF A CORONAL BRIGHT POINT

HUI TIAN,^{1,2} WERNER CURDT,¹ ECKART MARSCH,¹ AND JIANSAN HE²

Received 2008 April 18; accepted 2008 May 28; published 2008 June 24

ABSTRACT

We performed a systematic study of the Doppler shifts and electron densities measured in an EUV bright point (hereafter BP) observed in more than 10 EUV lines with formation temperatures from $\log(T/K) = 4.5$ to 6.3. Those parts of a BP seen in transition region and coronal lines are defined as its cool and hot components, respectively. We find that the transition from cool to hot occurs at a temperature around $\log(T/K) = 5.7$. The two components of the BP reveal a totally different orientation and Doppler-shift pattern, which might result from a twist of the associated magnetic loop system. The analysis of magnetic field evolution and topology seems to favor a two-stage heating process, in which magnetic cancellation and separator reconnection are powering, respectively, the cool and hot components of the BP. We also found that the electron densities of both components of the BP are higher than those of the surrounding quiet Sun, and comparable to or smaller than active region densities.

Subject headings: Sun: corona — Sun: magnetic fields — Sun: transition region — Sun: UV radiation

Online material: color figures

1. INTRODUCTION

Coronal bright points (BPs) are small-scale phenomena in the solar corona and characterized by enhanced emission in X-ray, extreme-ultraviolet (EUV), and radio wavelengths. They were found to be located at the network boundaries, where the quiet-Sun magnetic field is mainly concentrated (Habbal et al. 1990), and to expand with height (Gabriel 1976; Tian et al. 2008). Coronal BPs are typically 30"–40" in size, often with a bright core of 5"–10" (Madjarska et al. 2003). It is believed that a BP consists of several miniature dynamic loops (Sheeley & Golub 1979). The average lifetime of BPs is 20 hr in EUV (Zhang et al. 2001) and 8 hr in X-ray observations (Golub et al. 1974).

Madjarska et al. (2003) found that the Doppler shift of S VI in a BP is in the range of -10 to 10 km s⁻¹. Xia et al. (2003) found that the BPs in a coronal hole corresponded to a small blueshift of Ne VIII. Recently, both up- and downflows were detected in a BP (Brosius et al. 2007). The electron densities of BPs have been derived by using line pairs with a formation temperature of $\log(T/K) = 6.1$ – 6.3 (Ugarte-Urra et al. 2005; Brosius et al. 2008).

The evolution of a BP is strongly related to the underlying bipolar magnetic field (Webb et al. 1993; Falconer et al. 1998; Brown et al. 2001; Madjarska et al. 2003; Tian et al. 2007). Most BPs are more likely associated with the cancellation of magnetic features than their emergence (Webb et al. 1993). The energization of BPs may result from the interaction between two magnetic fragments of opposite polarities (Priest et al. 1994; Parnell et al. 1994; Von Rekowski et al. 2006), magnetic reconnection along separator field lines (Longcope 1998), or current sheets induced by photospheric motions (Büchner 2006; Santos & Büchner 2007).

Here we study the Doppler shifts and electron densities of a BP, which was observed by spectrometers on board *SOHO* and *Hinode* in a wide temperature range. We show the differences in the morphology and Doppler pattern between the cool

and hot components of this BP, and discuss possible mechanisms powering the radiation of the two components.

2. DATA ANALYSIS

The data set analyzed here was obtained by the Solar Ultraviolet Measurements of Emitted Radiation instrument (SUMER) (Wilhelm et al. 1995; Lemaire et al. 1997) and the EUV Imaging Spectrometer (EIS) (Culhane et al. 2007) on 2007 April 5. From 01:44 to 05:13 UTC, SUMER scanned three times a quiet-Sun region around disk center with a size of about 58" × 120", with each scan using different spectral windows. Slit 4 (1" × 120") was used and the exposure time was 90 s. The raster increment was about 1.5". EIS scanned from 03:15 to 04:09 UTC almost the same region by using the 1" slit with an exposure time of 90 s. The raster increment was about 2". The co-alignment between different SUMER images were done by using the cross-correlation technique. The He II (256.32 Å) image taken by EIS was used to co-align SUMER and EIS. During this period, an EUV BP was present in the field of view of the spectrometers. We selected some strong and clean spectral lines (see Table 1) to study the Doppler shifts, and some density-sensitive line pairs (see Fig. 4) to study the electron density of this BP. The values of rest wavelengths and formation temperatures were taken from Xia (2003) and the CHIANTI database (Dere et al. 1997; Landi et al. 2006).

We applied the standard procedures for correcting and calibrating the SUMER data, including decompression, flat-field correction, and detector corrections for geometrical distortion, local gain, and dead time. We also used the standard EIS calibrating procedures to subtract the dark current, remove the effects of cosmic rays and hot pixels, and make wavelength and absolute calibrations.

In order to build up intensity maps and Dopplergrams for the lines listed in Table 1, we applied a single Gaussian fit to each spectral profile. Then we estimated the line shift caused by the thermal deformation of the optical system of SUMER, using a similar method as described in Dammasch et al. (1999). Also, the fitted center of an EIS line was further corrected by taking into account the slit-tilt effect and orbital variation of the line position. We assumed a vanishing Doppler shift for

¹ Max-Planck-Institut für Sonnensystemforschung, 37191 Katlenburg-Lindau, Germany; tianhui924@gmail.com.

² Department of Geophysics, Peking University, 100871 Beijing, China.

TABLE 1
EMISSION LINES USED TO STUDY THE DOPPLER SHIFTS
OF THE BP

SUMER LINES			EIS LINES		
Ion	λ (Å)	$\log(T/K)$	Ion	λ (Å)	$\log(T/K)$
O II	833.332	4.5	Fe VIII	185.21	5.6
O III	833.749	4.9	Fe X	184.54	6.0
S IV	750.221	5.0	Fe XII	195.12	6.1
N IV	765.148	5.1	Fe XIII	202.04	6.2
O IV	787.711	5.2	Fe XIV	264.79	6.3
O V	758.677	5.4			
Ne VIII	770.428	5.8			

NOTE.—Here λ and T represent the rest wavelength and formation temperature, respectively.

each line when being averaged over the whole FOV. In this way we could calibrate the wavelength and obtained a Dopplergram for each line. Figures 1 and 2 show part of the resulting intensity maps and Dopplergrams, respectively.

During this time period, only the 96 minute MDI (Michelson Doppler Imager) magnetograms (Scherrer et al. 1995) were available. Figure 3 shows four magnetograms of the region in the vicinity of the BP taken at four different times. They were co-aligned by using the SUMER continuum intensity image obtained around 750 Å. The rightmost panel of Figure 3 is the magnetic skeleton of the BP region projected onto the plane of the photosphere. We first used the YAFTA (Yet Another Feature Tracking Algorithm) software (Welsch et al. 2004; DeForest et al. 2007) to select all individual magnetic concentrations and then reduced them to point sources. With the help of the MPOLE (the Magnetic Charge Topology and Minimum Current Corona model) software (Longcope & Klapper 2002), we could reconstruct the potential magnetic field topology produced by the set of point sources.

Some lines listed in Figure 4 are weak or blended. However, by using the method of single or double Gaussian fitting, we can still obtain a reliable intensity for each line in the BP region. Each of the density-sensitive line pairs was observed simultaneously. The theoretical relations between intensity ratios of line pairs and electron densities are shown in the top panel of Figure 4 and were taken from the CHIANTI database (Dere et al. 1997; Landi et al. 2006). Some blends discussed in Young et al. (2007) were also considered. We calculated intensities

and densities for each individual pixel within the corresponding contoured region in Figure 2 and then averaged those densities. We should mention that the O IV line pair was observed at about 11:40, when the BP was still there. The results are shown in the bottom of Figure 4.

3. RESULTS AND DISCUSSION

The temperature dependence of the emission of BPs and their different plasma properties at different wavelengths are not yet well understood. Most of the previous studies concentrated on BPs as seen in only one wavelength or band. Few authors have analyzed a BP seen simultaneously at different temperatures (Habbal & Withbroe 1981; Habbal et al. 1990; Habbal & Grace 1991; Ugarte-Urra et al. 2004; McIntosh 2007; Brosius et al. 2007).

Our Figure 1 reveals the rich morphology of a BP when observed with such a wide temperature coverage. The Dopplergrams and related intensity contours are shown in Figure 2. It is believed that BPs can experience variations in emission intensity and Doppler shift on a timescale of minutes (Habbal et al. 1990; Madjarska et al. 2003). However, the general emission pattern of some BPs can remain almost the same for 1–3 hr (see Fig. 2 in Brown et al. 2001). We should not exclude the possibility of the presence of a more or less stable component of emission and Doppler flow in some BPs. In our case, the orientation and Doppler pattern (especially the sharp boundary between patches of red- and blueshift) of the BP are both different at higher [$\log(T/K) > 5.7$] and lower [$\log(T/K) < 5.7$] temperatures. At lower temperatures, although the BP was observed at different times from 02:19 to 04:05, its orientation and Doppler pattern seem to be consistent and do not change too much.

Recently, Brosius et al. (2007) found Doppler shifts on opposite sides of a BP ranging from -15 to $+15$ km s $^{-1}$ in He II, and from -35 to $+35$ km s $^{-1}$ in Fe XVI. Our results demonstrate that the Doppler shifts in the BP are quite different in different lines. Patches of red- and blueshift are both found in the BP with comparable sizes. The absolute shift is largest in middle-transition region lines [$\log(T/K) = 4.9$ – 5.2] and can reach more than 10 km s $^{-1}$.

The most interesting phenomenon visible in Figure 2 is that




FIG. 1.—Intensity maps (on a logarithmic scale) of the lines listed in Table 1. The approximate time when the BP was scanned is shown in the lower right corner of each map. [See the electronic edition of the Journal for a color version of this figure.]




FIG. 2.—Dopplergrams of the lines listed in Table 1. The approximate time when the BP was scanned is shown in the lower right corner of each map. The black contours outline the positions of the BP as seen in different wavelengths. [See the electronic edition of the *Journal* for a color version of this figure.]

the boundary between up- and downflows in the BP is totally different, i.e., almost perpendicular, for lines with lower and higher temperatures. This boundary in Ne VIII resembles those of lines with a higher temperature. But the Doppler pattern of Ne VIII is similar to those of lines with a lower temperature in the surrounding quiet Sun. McIntosh (2007) classified the BPs seen in He II (304 Å) and soft X-ray as cool and hot BPs, respectively. Similarly, we define here the bright feature seen at coronal temperature as the hot component, and the corresponding bright emission at transition region temperature as the cool component of a BP. Considering the morphology and Doppler pattern of the BP, we found that a transition from the cool to the hot component of the BP occurs at a temperature of about $\log(T/K) = 5.7$.

The different boundary between up- and downflows at lower and higher temperatures in the BP might be a result of a syphon flow along a twisted loop system, which twists or spirals at its upper segment. Then the flow may lead to comparable red and blue shifts in the opposite legs, and result in a different boundary of the up- and downflows located between the lower and upper parts of the loop. In Figure 3, we found that the main negative magnetic source rotated relative to the positive source in a right-hand sense during the time period from 00:03 to 09:39. This movement might indicate a continual twist of the entire loop system. We have applied the force-free magnetic field extrapolation technique (Seehafer 1978) to reconstruct the 3D magnetic field in our BP region, but we could not reconstruct such a helical loop system. This does not mean that the twisted

loop is not there, because the BP may simply not be a force-free structure, and consequently the extrapolation used may not represent the real magnetic field adequately. However, the up- and downflows in our case might also be associated with magnetic reconnection, as discussed in Brosius et al. (2007).

The different emission morphology and Doppler pattern between the cool and hot components of the BP may imply a different powering mechanism of the two components. McIntosh (2007) proposed a two-stage heating process, in which magnetoconvection-driven reconnection occurs in and supplies energy to the cool BPs, whereupon the increased energy supply leads to an expansion of the loop system, which interacts with the overlying coronal magnetic flux through fast separator reconnection and produces hot BPs. From magnetograms taken before and after the observation periods of SUMER and EIS, we find that the main negative source in the left panels of Figure 3 is seen to approach and partly cancel the positive source. This cancellation might have started before and still be in process after our spectroscopic observations. We also find that the three separators (A4-B1, A4-B2, A4-B3) in the magnetic skeleton at 04:51 partly fit the orientation of the hot component of the BP. Thus, our observations seem to support the two-stage powering mechanism.

The electron density is vital to determine the radiative losses in coronal heating models. Here we derived separately the densities of the cool and hot components of the same BP. The results are summarized in Figure 4. Our derived values of the electron density in the range from $\log(T/K) = 6.1$ to 6.3 are




FIG. 3.—Left to right: MDI magnetograms taken at 00:03, 04:51, 09:39, and 19:12, and the magnetic skeleton at 04:51. The FOV corresponds to an x -coordinate range of $80''$ – $115''$ and a y -coordinate range of $5''$ – $37''$. In the magnetic skeleton, positive/negative magnetic point sources are labeled P/N and marked with plus/cross symbols; positive/negative nulls are labeled B/A and marked with triangles pointing down/up; spines and separators are shown as dotted and solid lines, respectively; dashed lines indicate where separatrix surfaces intersect the photosphere.




FIG. 4.—*Top*: Theoretical relation between the intensity ratios of six line pairs and the corresponding electron densities. *Bottom*: Derived electron densities and their uncertainties indicated by standard deviation bars.

similar to those of Ugarte-Urra et al. (2005) and Brosius et al. (2008). These densities are in the range of values typical of ARs, and slightly larger than quiet-Sun densities. The result of the higher value obtained with Fe XII compared to Si X is nothing new and a related discussion can be found in Ugarte-Urra et al. (2005).

The densities of the cool component of the BP have large uncertainties. But they can still be considered to be much higher than the density of the normal quiet Sun, which has an upper limit of $\log(N_e/\text{cm}^{-3}) = 9.87$ at $\log(T/\text{K}) = 5.25$ in Griffiths et al. (1999). Tripathi et al. (2008) derived a density of $\log(N_e/\text{cm}^{-3}) = 10.0\text{--}10.5$ at $\log(T/\text{K}) = 5.8\text{--}6.1$ inside the moss region of an AR. If we assume the density decreases with increasing temperature in the transition region, we can expect an even higher density of the AR at $\log(T/\text{K}) = 5.2\text{--}5.4$. It should be comparable to, or larger than, our derived densities at $\log(T/\text{K}) = 5.2\text{--}5.4$. So, the conclusion in Ugarte-Urra et al. (2005), namely that the BP plasma has more similarities to active-region than quiet-Sun plasma, may also be true for the cool component.

SUMER and MDI are instruments on board *SOHO*, an ESA and NASA mission. The SUMER project is financially supported by DLR, CNES, NASA, and the ESA PRODEX program (Swiss contribution). EIS is an instrument on board *Hinode*, a Japanese mission developed and launched by ISAS/JAXA, with NAOJ as domestic partner and NASA and STFC (UK) as international partners. It is operated by these agencies in cooperation with ESA and NSC (Norway). The work of Hui Tian and Jiansen He's team in PKU is supported by the National Natural Science Foundation of China under contracts 40574078 and 40436015. Hui Tian is now supported by China Scholarship Council for his stay in Germany. We thank the anonymous referee and M. Madjarska for helpful comments.

REFERENCES

- Brosius, J. W., Rabin, D. M., & Thomas, R. J. 2007, *ApJ*, 656, L41
 Brosius, J. W., Rabin, D. M., Thomas, R. J., & Landi, E. 2008, *ApJ*, 677, 781
 Brown, D. S., et al. 2001, *Sol. Phys.*, 201, 305
 Büchner, J. 2006, *Space Sci. Rev.*, 122, 149
 Culhane, J. L., et al. 2007, *Sol. Phys.*, 243, 19
 Dammasch, I. E., Wilhelm, K., Curdt, W., & Hassler, D. M. 1999, *A&A*, 346, 285
 DeForest, C. E., et al. 2007, *ApJ*, 666, 576
 Dere, K. P., et al. 1997, *A&AS*, 125, 149
 Falconer, D. A., Moore, R. L., Porter, J. G., & Hathaway, D. H. 1998, *ApJ*, 501, 386
 Gabriel, A. H. 1976, *Philos. Trans. R. Soc. London A*, 281, 339
 Golub, L., et al. 1974, *ApJ*, 189, L93
 Griffiths, N. W., Fisher, G. H., Woods, D. T., & Siegmund, O. H. W. 1999, *ApJ*, 512, 992
 Habbal, S. R., Dowdy, J. F., Jr., & Withbroe, G. L. 1990, *ApJ*, 352, 333
 Habbal, S. R., & Grace, E. 1991, *ApJ*, 382, 667
 Habbal, S. R., & Withbroe, G. L. 1981, *Sol. Phys.*, 69, 77
 Landi, E., et al. 2006, *ApJS*, 162, 261
 Lemaire, P., et al. 1997, *Sol. Phys.*, 170, 105
 Longcope, D. W. 1998, *ApJ*, 507, 433
 Longcope, D. W., & Klapper, I. 2002, *ApJ*, 579, 468
 Madjarska, M. S., Doyle, J. G., Teriaca, L., & Banerjee, D. 2003, *A&A*, 398, 775
 McIntosh, S. W. 2007, *ApJ*, 670, 1401
 Parnell, C. E., Priest, E. R., & Titov, V. S. 1994, *Sol. Phys.*, 153, 217
 Priest, E. R., Parnell, C. E., & Martin, S. F. 1994, *ApJ*, 427, 459
 Santos, J. C., & Büchner, J. 2007, *Ap&SS Trans.*, 3, 29
 Scherrer, P. H., et al. 1995, *Sol. Phys.*, 162, 129
 Seehafer, N. 1978, *Sol. Phys.*, 58, 215
 Sheeley, N. R., Jr., & Golub, L. 1979, *Sol. Phys.*, 63, 119
 Tian, H., Marsch, E., Tu, C.-Y., Xia, L.-D., & He, J.-S. 2008, *A&A*, 482, 267
 Tian, H., Tu, C.-Y., He, J.-S., & Marsch, E. 2007, *Adv. Space Res.*, 39, 1853
 Tripathi, D., Mason, H. E., Young, P. R., & Del Zanna, G. 2008, *A&A*, 481, L53
 Ugarte-Urra, I., Doyle, J. G., & Del Zanna, G. 2005, *A&A*, 435, 1169
 Ugarte-Urra, I., Doyle, J. G., Madjarska, M. S., & O'Shea, E. 2004, *A&A*, 418, 313
 Von Rekowski, B., Parnell, C. E., & Priest, E. R. 2006, *MNRAS*, 366, 125
 Webb, D. F., Martin, S. F., Moses, D., & Harvey, J. W. 1993, *Sol. Phys.*, 144, 15
 Welsch, B. T., Fisher, G. H., Abbett, W. P., & Regnier, S. 2004, *ApJ*, 610, 1148
 Wilhelm, K., et al. 1995, *Sol. Phys.*, 162, 189
 Xia, L.-D. 2003, Ph.D. thesis, Georg-August-Univ.
 Xia, L. D., Marsch, E., & Curdt, W. 2003, *A&A*, 399, L5
 Young, P. R., et al. 2007, *PASJ*, 59, S857
 Zhang, J., Kundu, M. R., & White, S. M. 2001, *Sol. Phys.*, 198, 347